



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07D 401/12, A61K 31/44</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 92/05170</b> <b>(43) International Publication Date:</b> 2 April 1992 (02.04.92)
<b>(21) International Application Number:</b> PCT/GB91/01553 <b>(22) International Filing Date:</b> 11 September 1991 (11.09.91)  <b>(30) Priority data:</b> 9020030.4           13 September 1990 (13.09.90)   GB 9106079.8           22 March 1991 (22.03.91)       GB 9106092.1           22 March 1991 (22.03.91)       GB 9106094.7           22 March 1991 (22.03.91)       GB  <b>(71) Applicant (for all designated States except US):</b> BEECHAM GROUP PLC [GB/GB]; SB House, Great West Road, Brentford, Middlesex TW8 9BD (GB).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only) :</b> FORBES, Ian, Thomson [GB/GB]; MARTIN, Roger, Thomas [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB).		<b>(74) Agent:</b> VALENTINE, Jill; SmithKline Beecham, Corporate Patents, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ (GB).  <b>(81) Designated States:</b> AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent), US.  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> INDOLE UREAS AS 5 HT RECEPTOR ANTAGONIST  <b>(57) Abstract</b>  Indole ureas, pharmaceutical compositions containing them and processes for their preparation.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU <sup>+</sup>	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE*	Germany	MC	Monaco	US	United States of America
DK	Denmark				

<sup>+</sup> Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.


## INDOLE UREAS AS 5 HT RECEPTOR ANTAGONIST

This invention relates to compounds having pharmacological activity, to a process for their preparation, to compositions containing them and to their use in the treatment of mammals.

P. Fludzinski *et. al.*, J. Med. Chem. 1986 29 2415-2418 describes N-(1,2-dimethyl-3-ethyl-1H-indol-5-yl)-N'-(3-trifluoromethylphenyl)urea which shows selectivity for the rat stomach fundus serotonin receptor.

A class of compounds has now been discovered, which compounds have been found to have 5HT<sub>1C</sub> receptor antagonist activity. 5HT<sub>1C</sub> receptor antagonists are believed to be of potential use in the treatment of CNS disorders such as anxiety, depression, obsessive compulsive disorders, migraine, anorexia, Alzheimers disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse and/or schizophrenia.

Accordingly, the present invention provides a compound of formula (I) or a pharmaceutically acceptable salt thereof:



wherein:

-2-

R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are independently hydrogen or C<sub>1-6</sub> alkyl;  
R<sub>4</sub> is hydrogen, C<sub>1-6</sub> alkyl, halogen, hydroxy or NR<sub>8</sub>R<sub>9</sub> where  
R<sub>8</sub> and R<sub>9</sub> are independently hydrogen or C<sub>1-6</sub> alkyl;  
R<sub>5</sub> and R<sub>6</sub> are independently hydrogen or C<sub>1-6</sub> alkyl; and  
5 R<sub>7</sub> is hydrogen, C<sub>1-6</sub> alkyl or halogen; and wherein the urea  
moiety is attached at the 4-, 5- or 6-position of the indole  
ring.

Alkyl moieties within the variables R<sub>1</sub> to R<sub>9</sub> are preferably  
10 C<sub>1-3</sub> alkyl, such as methyl, ethyl, n- and iso- propyl, most  
preferably methyl, ethyl and n-propyl.

Suitable R<sub>4</sub> and R<sub>7</sub> halogens include chloro and bromo.

15 Examples of R<sub>1</sub> include hydrogen, methyl, ethyl and n-propyl,  
preferably methyl. R<sub>2</sub> is preferably methyl or hydrogen and  
R<sub>3</sub> is hydrogen, methyl, ethyl, n-propyl, iso-propyl or  
n-hexyl.

20 Preferably R<sub>4</sub> is hydrogen, chloro, hydroxy or dimethylamino,  
most preferably hydrogen.

Preferably R<sub>5</sub>, R<sub>6</sub> and R<sub>7</sub> are independently hydrogen or  
methyl.

25

The urea moiety may be attached at the 2-, 3-, 4-, 5- or 6-  
position of the pyridine ring, preferably the 3-, 4- or  
5-position.

30 The urea moiety is preferably attached at the 4- or  
5-position of the indole ring.

The compounds of the formula (I) can form acid addition  
salts with acids, such as conventional pharmaceutically  
35 acceptable acids, for example maleic, hydrochloric,  
hydrobromic, phosphoric, acetic, fumaric, salicylic, citric,

-3-

lactic, mandelic, tartaric and methanesulphonic.

Compounds of formula (I) may also form solvates such as hydrates, and the invention also extends to these forms.


5 When referred to herein, it is understood that the term 'compound of formula (I)' also includes solvates thereof.

When  $R_5$  and/or  $R_6$  are hydrogen or when  $R_4$  is 2- or 4-hydroxy or  $NR_8R_9$  and at least one of  $R_8$  and  $R_9$  are hydrogen the  
10 compounds of formula (I) may exist tautomerically in more than one form. The invention extends to each of these forms and mixtures thereof.

Certain compounds of formula (I) are capable of existing in  
15 stereoisomeric forms including enantiomers and the invention extends to each of these stereoisomeric forms and to mixtures thereof including racemates. The different stereoisomeric forms may be separated one from the other by the usual methods, or any given isomer may be obtained by  
20 stereospecific or asymmetric synthesis.

The present invention also provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process  
25 comprises

(a) the coupling of a compound of formula (II);




(II)

35

-4-

with a compound of formula (III);




10 wherein B is attached at the 4-, 5- or 6-position of the indole ring and A and B contain the appropriate functional group(s) necessary to form the moiety  $-NR_5'CONR_6'-$  when coupled, wherein  $R_5'$  and  $R_6'$  are  $R_5$  and  $R_6$  as defined in formula (I) or groups convertible thereto, and the variables

15  $R_1'$ ,  $R_2'$ ,  $R_3'$ ,  $R_4'$  and  $R_7'$  are  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  and  $R_7$  respectively, as defined in formula (I), or groups convertible thereto, and thereafter optionally and as necessary and in any appropriate order, converting any  $R_1'$ ,  $R_2'$ ,  $R_3'$ ,  $R_4'$ ,  $R_5'$ ,  $R_6'$  and  $R_7'$  when other than  $R_1$ ,  $R_2$ ,  $R_3$ ,

20  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$  respectively to  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$ , interconverting  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$ , and forming a pharmaceutically acceptable salt thereof, or

(b) cyclising a compound of formula (IV):

25



wherein  $R_4'$ ,  $R_5'$ ,  $R_6'$  and  $R_7'$  are as defined in formulae (II) and (III) and C and D contain the appropriate

35 functional group(s) necessary to form the indole ring

-5-

substituted by  $R_1'$ ,  $R_2'$  and  $R_3'$  as defined in formula (III), and thereafter optionally and as necessary in any appropriate order, converting any  $R_1'$ ,  $R_2'$ ,  $R_3'$ ,  $R_4'$ ,  $R_5'$ ,  $R_6'$  and  $R_7'$  when other than  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$ , to  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$ , interconverting  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$  and forming a pharmaceutically acceptable salt.

Suitable examples of groups A and B are

10

- (i) A is  $-N=C=O$  and B is  $-NHR_6'$ ,
- (ii) A is  $-NHR_5'$  and B is  $-N=C=O$ ,
- (iii) A is  $-NR_5'COL$  and B is  $-NHR_6'$ ,
- (iv) A is  $-NHR_5'$  and B is  $-NR_6'COL$ , or

15 (v) A is halogen and B is  $-NR_6'CONHR_5'$ ,

wherein  $R_5'$  and  $R_6'$  are as defined above and L is a leaving group. Examples of suitable leaving groups L include halogen such as chloro or bromo, imidazole, or phenoxy or phenylthio optionally substituted for example with halogen.

20

When A is  $-N=C=O$  and B is  $NHR_6'$  or when A is  $NHR_5'$  and B is  $-N=C=O$  the reaction is suitably carried out in an inert solvent for example dichloromethane or toluene at ambient temperature.

25

When A is  $-NR_5'COL$  and B is  $-NHR_6'$  or when A is  $-NHR_5'$  and B is  $-NR_6'COL$ , the reaction is suitably carried out in an inert solvent such as dichloromethane at ambient temperature optionally in the presence of a base, such as triethylamine  
30 or in dimethylformamide at ambient or elevated temperature.

When A is halogen and B is  $-NR_6'CONHR_5'$ , the reaction is suitably carried out in an inert solvent such as toluene at elevated temperature, optionally in the presence of a base.

35

-6-

The cyclisation of the compound of formula (IV) may be effected using standard methodology such as described in Comprehensive Heterocyclic Chemistry 1984 4, 313 et seq. or J. Het. Chem. 1988 25 p.1 et seq.

5

Examples of the more important routes include the Leimgruber synthesis, the Fischer synthesis and the Japp-Klingemann variation and the Madelung synthesis.

10 Examples of the groups C and D thus include

(vi)  $C = NO_2$  and  $D = CH=CH-NZ_2$  where each Z is independently  $C_{1-6}$  alkyl or together represent  $C_{2-7}$  alkylene;

15

(vii)  $C = NR_1'-N=C(R_2')-CH_2R_3'$  and  $D = H$ ;

(viii)  $C = NH-N=C(CO_2X)-CH_2R_3'$  and  $D = H$  where X is  $C_{1-6}$  alkyl; and

20

(ix)  $C = NR_1'COR_2'$  and  $D = CH_2R_3'$ .

In reaction variant (vi) (Leimgruber synthesis) the compound of formula (IV) is prepared from the 2-methylnitrophenyl  
25 urea by treatment with a dialkylacetal of the dialkylformamide  $OHCNZ_2$  with heating and the product of formula (IV) cyclised by hydrogenation over a suitable catalyst such as palladium and charcoal optionally under pressure to yield the compound of formula (I) where  
30  $R_1=R_2=R_3=H$ .

In reaction variant (vii) (Fischer synthesis) the compound of formula (IV) is prepared from the hydrazinophenyl urea by dehydration, preferably by heating, with the appropriate  
35 ketone  $R_2'COCH_2R_3'$  and the product of formula (IV) cyclised by heating with an acid catalyst such as hydrochloric or sulphuric acid.



-7-

In reaction variant (viii) (Japp-Klingemann synthesis) the compound of formula (IV) is prepared from the aminophenyl urea by diazotisation followed by treatment for example with  $\text{CH}_3\text{COCH}(\text{CO}_2\text{X})-\text{CH}_2\text{R}_3'$  where X is  $\text{C}_{1-6}$  alkyl under basic conditions in aqueous alcohol as solvent.

The product of formula (IV) may then be cyclised as in the Fischer synthesis above.

10 In reaction variant (ix) (Madelung synthesis) the compound of formula (IV) is cyclised with base in an inert solvent optionally with heating.

Suitable examples of groups  $\text{R}_2'$ ,  $\text{R}_3'$ ,  $\text{R}_4'$ , and  $\text{R}_7'$  which are convertible to  $\text{R}_2$ ,  $\text{R}_3$ ,  $\text{R}_4$ , and  $\text{R}_7$  respectively, include acyl groups which are introduced conventionally and may be converted to the corresponding alkyl group by conventional reduction, such as using sodium borohydride in an inert solvent followed by hydrogenolysis in an inert solvent and alkoxy carbonyl groups which may be converted to hydrogen by hydrolysis and decarboxylation. When  $\text{R}_4$  is hydroxy it is preferably protected in the compound of formula (II) as, for example, an aryloxy group such as benzyloxy which is removed by hydrogenation.

25

Suitable examples of a group  $\text{R}_1'$  which is convertible to  $\text{R}_1$ , include typical N-protecting groups such as alkoxy carbonyl, in particular t-butyloxy carbonyl, acetyl, trifluoroacetyl, benzyl and para-methoxybenzyl which are converted to  $\text{R}_1$  hydrogen using conventional conditions.

Suitable examples of groups  $\text{R}_5'$  and  $\text{R}_6'$  which are convertible to  $\text{R}_5$  and  $\text{R}_6$  respectively include alkoxy carbonyl and benzyl or para-methoxybenzyl which are converted to  $\text{R}_5$  and/or  $\text{R}_6$  hydrogen using conventional conditions.

35

-8-

Interconversions of  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$  are carried out by conventional procedures. For example, in the case wherein  $R_1$ ,  $R_2$  and  $R_3$  are  $C_{1-6}$  alkyl and  $R_5$  and  $R_6$  are hydrogen it is possible to introduce a  $C_{1-6}$  alkyl group at both the  $R_5$  and  $R_6$  positions by conventional alkylation using 2 molar equivalents of a  $C_{1-6}$  alkyl halide and 2 molar equivalents of a suitable base in an inert solvent. Monoalkylation can be achieved using 1 molar equivalent of a  $C_{1-6}$  alkyl halide and base using conventional conditions.  $R_1$   $C_{1-6}$  alkyl groups may also be introduced by conventional alkylation, for example using a  $C_{1-6}$  alkyl halide and base such as sodium hydride.

$R_4$  halo and  $R_7$  halo may be introduced by selective halogenation of the pyridine ring or indole ring respectively using conventional conditions.

It should be appreciated that it may be necessary to protect any  $R_1$  to  $R_7$  hydrogen variables which are not required to be interconverted.

Suitable protecting groups and methods for their attachment and removal are conventional in the art of organic chemistry, such as those described in Greene T.W. 'Protective groups in organic synthesis' New York, Wiley (1981).

It is preferable, however, to introduce and interconvert the groups  $R_1$  to  $R_7$  before coupling compounds of formulae (II) and (III) together, or cyclising the compound of formula (IV).

Compounds of formula (II) in which A is  $NHR_5'$  are known compounds or can be prepared analogously to known compounds. For example, the compounds of formula (II) in which A is 3-amino and  $R_4'$  is hydrogen, 2-chloro or 6-chloro, and A is

-9-

2-amino and  $R_4'$  is 3-benzyloxy are commercially available from the Aldrich Chemical Company in the UK.  $R_5'$   $C_{1-6}$  alkyl groups may be introduced conventionally, for example by reductive alkylation or acylation and reduction.

5

Compounds of formula (II) in which A is  $-N=C=O$  may be prepared by treating a compound of formula (II) in which :

i) A is amino, with phosgene or a phosgene equivalent, in  
10 the presence of excess base in an inert solvent.

ii) A is acylazide (i.e.  $CON_3$ ), via the nitrene, by thermal rearrangement using conventional conditions (ref L.S. Trifonov et al, *Helv. Chim. Acta* 1987 70 262).

15

iii) A is  $CONH_2$ , via the nitrene intermediate using conventional conditions.

Compounds of formula (II) in which A is  $-NR_5'COL$  may be  
20 prepared by reacting a compound of formula (II) in which A is  $-NHR_5'$  with phosgene or a phosgene equivalent, in an inert solvent, at low temperature, if necessary in the presence of one equivalent of a base such as triethylamine.

25 Compounds of formula (II) in which A is halogen and  $R_4'$  is hydrogen are commercially available.

Compounds of formula (III) in which B is  $NHR_6'$  are known compounds or can be prepared analogously to known compounds,  
30 for example by reduction of the corresponding nitroindole by catalytic hydrogenation over Pd/C by the method of P.

Fludzinski et al *J. Med. Chem.*, 1986, 29 2415.

Specifically, the compound of formula (III) in which  $R_1'$  and  $R_2'$  are methyl,  $R_3'$  is ethyl,  $R_6'$  and  $R_7'$  are hydrogen and B

-10-

is  $\text{NH}_2$  is prepared using a procedure similar to that described by Fludzinski.

The nitroindoles are commercially available, for example 5-nitroindole, or may be prepared conventionally (Comprehensive Heterocyclic Chemistry Vol. 4 p. 313 et. seq. (Pergamon Press 1984) and J. Het. Chem. 1988 25 p.1 et. seq.)

10 An  $\text{R}_2'$  alkoxycarbonyl group may be eliminated to give  $\text{R}_2'$  hydrogen, generally under the conditions effecting formation of the nitroindole or as a subsequent step in the process.

$\text{R}_6'$  alkyl groups may be introduced conventionally, for example by reductive alkylation or acylation and reduction.  $\text{R}_7'$   $\text{C}_{1-6}$  alkyl groups may be introduced ortho to a nitro substituent by alkylation using a procedure similar to that described in G. Bartoli et. al., J. Org. Chem. 1986 51 3694 and Tetrahedron 1987 43 4221.

20

Compounds of formula (III) in which B is  $-\text{N}=\text{C}=\text{O}$  may be prepared by treating a compound of formula (III) in which :

i) B is amino, with phosgene or a phosgene equivalent, in the presence of excess base in an inert solvent.

ii) B is acylazide (i.e.  $\text{CON}_3$ ), via the nitrene, by thermal rearrangement using conventional conditions.

30 iii) B is  $\text{CONH}_2$ , via the nitrene intermediate using conventional conditions.

Compounds of formula (III) in which B is  $-\text{NR}_6'\text{COL}$  may be prepared by reacting a compound of formula (III) in which B is  $-\text{NHR}_6'$  with phosgene or a phosgene equivalent, in an inert solvent, at low temperature, if necessary in the

-11-

presence of one equivalent of a base such as triethylamine.

Compounds of formula (III) in which B is  $-NR_6'CONHR_5'$  can be prepared from the corresponding precursor where B is  $NHR_6'$  by reaction with an  $R_5'$  isocyanate under conventional conditions.

Examples of phosgene equivalents include triphosgene, carbonyldiimidazole, phenyl chloroformate and phenyl chlorothioformate.

Novel intermediates of formula (III) also form part of the invention.

Compounds of formula (IV) may be prepared from the appropriate aminophenyl derivative analogously to compounds of formula (I). Intermediates of formula (IV) also form part of the invention.

Pharmaceutically acceptable salts may be prepared conventionally by reaction with the appropriate acid or acid derivative.

Compounds of formula (I) and their pharmaceutically acceptable salts have  $5HT_{1C}$  receptor antagonist activity and are believed to be of potential use in the treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse and/or schizophrenia. Thus the invention also provides a compound of formula (I) or a pharmaceutically acceptable salt thereof, for use as a therapeutic substance, in particular in the treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia,

-12-

panic attacks, withdrawal from drug abuse and/or schizophrenia.

The invention further provides a method of treatment or  
5 prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse and/or schizophrenia in mammals including humans, which comprises administering to the sufferer a  
10 therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

In another aspect, the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable  
15 salt thereof in the manufacture of a medicament for the treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse and/or schizophrenia.

20

The present invention also provides a pharmaceutical composition, which comprises a compound of formula (I) or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

25

A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral or rectal administration and, as such, may be in  
30 the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusable solutions or suspensions or suppositories. Orally administrable compositions are generally preferred.

35

Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as

-13-

binding agents, fillers, tableting lubricants, disintegrants and acceptable wetting agents. The tablets may be coated according to methods well known in normal pharmaceutical practice.

5

Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before  
10 use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous vehicles (which may include edible oils), preservatives, and, if desired, conventional flavourings or colourants.

15

For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle. The compound, depending on the vehicle and  
20 concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing.

Advantageously, adjuvants such as a local anaesthetic, 25 preservatives and buffering agents are dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is  
30 suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition  
35 to facilitate uniform distribution of the compound.

-14-

The composition may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration.

5 The dose of the compound used in the treatment of the  
aforementioned disorders will vary in the usual way with the  
seriousness of the disorders, the weight of the sufferer,  
and other similar factors. However, as a general guide  
suitable unit doses may be 0.05 to 1000 mg, more suitably  
10 0.05 to 20.0 mg, for example 0.2 to 5 mg; and such unit  
doses may be administered more than once a day, for example  
two or three a day, so that the total daily dosage is in the  
range of about 0.01 to 100 mg/kg; and such therapy may  
extend for a number of weeks or months.

15

When administered in accordance with the invention, no  
unacceptable toxicological effects are expected with the  
compounds of the invention.

20 The following Examples illustrate the preparation of  
pharmacologically active compounds of the invention. The  
following Descriptions illustrate the preparation of  
intermediates to compounds of the present invention.



-15-

Description 15-Amino-1,2-dimethyl-3-ethyl-1H-indole (D1)

5 The title compound was prepared using a procedure similar to that described by P. Fludzinski et al in J. Med. Chem., 1986, 29, 2415.

Description 2

10

1-Methyl-5-nitro-1H-indole (D2)

To a stirred suspension of sodium hydride (5.0g; 167 mM) in dimethylformamide (200 ml) at 0°C under nitrogen was added  
15 5-nitroindole (25g; 154 mM) in dimethylformamide. After stirring for 0.5h, iodomethane (10.5ml; 168mM) in dimethylformamide (50 ml) was added, and stirring was continued for 2h. The reaction mixture was then quenched with water, and poured onto excess water with stirring.  
20 Filtration afforded the title compound (27.4g; 94%).

NMR (CDCl<sub>3</sub>)  $\delta$ :

3.88 (3H, s), 6.68 (1H, d, J=3) 7.21 (1H, d, J=3) 7.34  
25 (1H, d, J=8) 8.13 (1H, dd, J=8, 2) 8.59 (1H, d, J=2).

Description 35-Amino-1-methyl-1H-indole (D3)

30

A mixture of the nitroindole (D2) (5g; 28.4 mM) and 5% palladium on charcoal in ethanol (300 ml) was hydrogenated at 60 p.s.i. ( $4.14 \times 10^5$  Pa) at room temperature for 3h. Removal of the catalyst by filtration followed by

-16-

evaporation of the solvent gave the title compound (3.39g; 95%).

NMR (CDCl<sub>3</sub>)  $\delta$ :

5

3.20 (2H, broad s), 3.70 (3H, s), 6.28 (1H, d, J=3)  
6.68 (1H, dd, J=8, 2), 6.92 (1H, d, J=2), 6.96 (1H, d, J=3),  
7.12 (1H, d, J=8)

10 Description 4

3-Pyridyl isocyanate (D4)

The title compound was prepared from 3-pyridinecarbonyl  
15 azide in toluene using a procedure similar to that described  
by L.S. Trifonov et al in Helv. Chim. Acta, 1987, 70, 262.

Description 5

20 5-Nitro-1,2,3-trimethyl-1H-indole (D5)

The title compound was prepared in 99% yield from 2,3-  
dimethyl-5-nitroindole using a procedure similar to that in  
Description 2.

25

NMR (CDCl<sub>3</sub>)  $\delta$ :

2.28 (3H, s), 2.38 (3H, s), 3.71 (3H, s), 7.22 (1H, d, J=8),  
8.06 (1H, dd, J=8, 2), 8.46 (1H, d, J=2).

30

-17-

Description 65-Amino-1,2,3-trimethyl-1H-indole (D6)

5 The title compound was prepared in 91% yield from the nitroindole (D5) using a procedure similar to that in Description 3.

NMR (CDCl<sub>3</sub>)  $\delta$ :

10

2.18 (3H, s), 2.29 (3H, s), 3.00 (2H broad s), 3.57 (3H, s),  
6.60 (1H, dd, J=8, 2) 6.80 (1H, d, J=2), 7.03 (1H, d, J=8).

Description 7

15

5-Nitro-1-propyl-1H-indole (D7)

The title compound was prepared in 96% yield from 5-nitroindole and propyl iodide using a procedure similar to  
20 that in Description 2.

NMR (CDCl<sub>3</sub>)  $\delta$ :

0.96 (3H, t, J=7), 1.90 (2H, h, J=7), 4.13 (2H, t, J=7),  
25 6.68 (1H, d, J=3), 7.26 (1H, d, J=3), 7.37 (1H, d, J=8),  
8.10 (1H, dd, J=8, 2), 8.59 (1H, d, J=2).

Description 830 5-Amino-1-propyl-1H-indole (D8)

The title compound was prepared in 100% yield from the nitroindole (D7) using a procedure similar to that in Description 3.

35

-18-

NMR (CDCl<sub>3</sub>)  $\delta$ :

0.91 (3H, t, J=7, 1.83 (2H, h, J=7), 3.38 (2H, broad s),  
4.02 (2H, t, J=7), 6.29 (1H, d, J=3), 6.68 (1H, dd, J=8, 2),  
5 6.93 (1H, d, J=2), 7.01 (1H, d, J=3), 7.14 (1H, d, J=8).

Description 91-Methyl-4-nitro-1H-indole (D9)

10

To a stirred suspension of sodium hydride (0.14g; 3.41 mM) in dimethylformamide (10 ml) at 0°C under nitrogen was added 4-nitroindole (0.5g; 3.1 mM) in dimethylformamide. After stirring for 0.5h, iodomethane (0.21 ml; 3.41 mM) in  
15 dimethylformamide (1 ml) was added, and stirring was continued for 1h. The reaction mixture was then quenched with water, and poured onto excess water with stirring. Filtration afforded the title compound (0.5g; 92%).

20 NMR (CDCl<sub>3</sub>)  $\delta$ :

3.89 (3H, s), 7.30 (3H, m), 7.66 (1H, d, J=8), 8.15 (1H, d, J=8).

25 Description 104-Amino-1-methyl-1H-indole (D10)

A mixture of the nitroindole (D9) (0.5g; 2.8 mM) and 5%  
30 palladium on charcoal in ethanol (75 ml) was hydrogenated at 60 p.s.i. ( $4.14 \times 10^5$  Pa) at room temperature for 2h. Removal of the catalyst by filtration followed by evaporation of the solvent gave the title compound (0.44g; 97%).

35

-19-

NMR (CDCl<sub>3</sub>)  $\delta$ :

3.76 (3H, s), 6.42 (1H, d, J=2), 6.45 (1H, d, J=8), 6.81  
(1H, d, J=8), 6.96 (1H, d, J=2), 7.05 (1H, t, J=8).

5

Description 111-Methyl-6-nitro-1H-indole (D11)

10 To a solution of sodium hydride (0.27g; 6.8mM) in  
dimethylformamide (4 ml) at 0°C under nitrogen, was added 6-  
nitroindole (1g; 6.2mM) in dimethylformamide (12 ml). After  
stirring at room temperature for 0.5h, iodomethane (0.42 ml;  
6.8 mM) in dimethylformamide (1 ml) was added and stirring  
15 continued for 1h. The reaction mixture was then quenched  
with water, and poured onto excess water with stirring.  
Filtration afforded the title compound (1.03g; 94%).

NMR (CDCl<sub>3</sub>)  $\delta$ :

20

3.60 (3H, s), 6.60 (1H, d, J=4), 7.35 (1H, d, J=4),  
7.55 (1H, d, J=10), 8.10 (1H, dd, J=10, 2), 8.34  
(1H, d, J=2)

25 Description 126-Amino-1-methyl-1H-indole (D12)

A mixture of the nitroindole (D11) (0.8g; 4.55 mM) and 5%  
30 palladium on charcoal in ethanol (150 ml) was hydrogenated  
at 60 p.s.i. ( $4.14 \times 10^5$  Pa) at room temperature for 2h.  
Removal of the catalyst by filtration followed by  
evaporation of the solvent gave the crude product.

-20-

Chromatography on silica using dichloromethane as eluant afforded the title compound (0.3g; 45%).

NMR (CDCl<sub>3</sub>)  $\delta$ :

5

3.68 (3H, s), 6.38 (1H, d, J=3), 6.55-6.65 (2H, m), 6.88 (1H, d, J=3), 7.40 (1H, s, J=10).

Description D13

10

3-Methylaminopyridine (D13)

A mixture of 3-aminopyridine (5.76g; 60 mM) in triethylorthoformate (49 ml) was refluxed with stirring for 15 5h. The excess solvent was removed in vacuo to give an oil (8.53g, 93%). The oil was dissolved in ethanol (30 ml) and cooled in ice. To this solution was added sodium borohydride (2.58g, 68.3 mM) portionwise and left to stir at room temperature for 17h. The solution was cooled in an ice 20 bath, and water added slowly (3 ml), followed by 5N HCl until no further evolution of gas was observed. The pH was adjusted to 7, then extracted using ethyl acetate, washed with water, dried and evaporated to give an oil (5.10g; 83%). Chromatography on silica using dichloromethane as 25 eluant afforded the title compound (1.43g; 22%).

NMR (CDCl<sub>3</sub>)  $\delta$ : 2.82 (3H, s), 4.12 (1H, s), 6.87 (1H, dd, J=8,3), 7.09 (1H, m), 7.95 (1H, dd, J=3,1), 8.02 (1H, d, J=3).

30

-21-

Description 14N-(1-Methyl-1H-indol-5-yl)formamide (D14)

5 To acetic anhydride (1.68 ml; 15 mM) at 0°C was added 98% formic acid (0.8 ml; 21 mM) dropwise under a nitrogen atmosphere, to generate acetic formic anhydride. The solution was heated at 50-60°C for 2h then cooled to room temperature. Dichloromethane (2 ml) was added and the  
10 solution was cooled to -20°C before adding a solution of aminoindole (D3) (1g, 6.88 mM) in dichloromethane (4 ml). The mixture was stirred at room temperature for 17h then evaporated to dryness to give a brown oil (1.34g) Chromatography on silica using ethyl acetate as eluant  
15 afforded the title compound (1.05g; 88%).

NMR (CDCl<sub>3</sub>) δ: Complex spectrum due to amide isomers

Found: M<sup>+</sup> 174

20 C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O requires 174

Description 151-Methyl-5-methylamino-1H-indole (D15)

25

To a suspension of lithium aluminium hydride (0.33g; 8.7 mM) in dry tetrahydrofuran (15 ml) at 0°C under a nitrogen atmosphere was added the amide (D14) (1.0g; 5.74 mM). The solution was left to stir at room temperature for 17h,  
30 cooled to 0°C and then water (3.5 ml), 5N sodium hydroxide solution (3.5 ml) and then water (5 ml) added in that order. The solution was left to stir for 10 min, then filtered and evaporated to give a brown oil (0.89g). Chromatography on silica using dichloromethane as eluant gave the title  
35 compound (0.57g; 62%).

-22-

NMR (CDCl<sub>3</sub>)  $\delta$ : 2.9 (3H, s), 3.52 (1H, s), 3.73 (3H, s),  
6.33 (1H, d, J=3), 6.69 (1H, dd, J=8,1),  
6.87 (1H, d, J=1), 6.97 (1H, d, J=3),  
7.16 (1H, d, J=8).

5

Description 161,4-Dimethyl-5-nitroindole (D16)

10 The title compound was prepared from 1-methyl-5-nitroindole (D2) using a procedure similar to that described by G.Bartoli *et al* in J.Org.Chem. 1986, 51, 3694 and Tetrahedron 1987, 43, 4221. This gave a yellow solid, m.p. 120-3°C, in 64% yield.

15

NMR (CDCl<sub>3</sub>)  $\delta$ : 2.84 (3H, s), 3.83 (3H, s), 6.71 (1H, d, J=3), 7.18 (1H, d, J=3), 7.20 (1H, d, J=8), 7.99 (1H, d, J=8).

20 Found: M<sup>+</sup> 190

C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub> requires 190

Found: C, 63.0; H, 5.3; N, 14.6%. C<sub>10</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub> requires C, 63.1; H, 5.3; N, 14.7%

25

Description 175-Amino-1,4-dimethylindole (D17)

30 The title compound was prepared from 1,4-dimethyl-5-nitroindole (D16) by catalytic hydrogenation as described in Description 3. This gave a dark purple oil in 92% yield.



-23-

NMR (CDCl<sub>3</sub>) δ: 2.34 (3H, s), 3.1 (2H, bs), 3.72 (3H, s), 6.37 (1H, d, J=3), 6.71 (1H, d, J=8), 6.98 (1H, d, J=3), 7.02 (1H, d, J=8).

5

Description 18

N-(1-Methyl-1H-indol-5-yl)-N'-(3-benzyloxy pyrid-2-yl)urea (D18)

10

The title compound was prepared from 5-amino-1-methyl-1H-indole (D3), carbonyl diimidazole and 2-amino-3-benzyloxy pyridine using a procedure similar to that described in Example 1.

15

NMR (D<sub>6</sub>-DMSO) δ: 3.80 (3H, s), 5.44 (2H, s), 6.40 (1H, d, J=6), 7.35 (7H, m), 7.61 (2H, dd, J=13,3), 7.80 (2H, d, J=3), 7.94 (2H, d, J=6).

20

Description 19

3-Ethyl-2-methyl-5-nitro-1H-indole (D19)

25 The title compound was prepared using a procedure identical to that described by P. Fludzinski et al in J.Med.Chem., 1986, 29, 2415.

Description 20

30

1,3-Diethyl-2-methyl-5-nitro-1H-indole (D20)

The title compound was prepared in 92% yield from the nitroindole (D19), sodium hydride, and iodoethane using a  
35 procedure similar to that described in Description 2.

-24-

NMR (CDCl<sub>3</sub>)  $\delta$ : 1.23 (3H, t, J=8), 1.36 (3H, t, J=8),  
2.40 (3H, s), 2.73 (2H, q, J=8), 4.12  
(2H, q, J=8), 7.21 (1H, d, J=9), 8.02  
(1H, dd, J=9, 2), 8.49 (1H, d, J=2).

5

Description 215-Amino-1,3-diethyl-2-methyl-1H-indole (D21)

10 The title compound was prepared in 86% yield from the  
nitroindole (D20) using a procedure similar to that in  
Description 3.

15 NMR (CDCl<sub>3</sub>)  $\delta$ : 1.22 (3H, t, J=8), 1.37 (3H, t, J=8),  
2.39 (3H, s), 2.75 (2H, q, J=8), 4.14  
(2H, q, J=8), 7.26 (1H, d, J=8), 8.04  
(1H, dd, J=8, 1), 8.50 (1H, d, J=1).

Description 22

20

2-Methyl-5-nitro-3-propyl-1H-indole (D22)

The title compound was prepared in 94% yield from the 4-  
nitrophenylhydrazone of 2-hexanone using the method of  
25 Fludzinski et al described in J.Med. Chem., 1986, 29, 2415.

30 NMR (CDCl<sub>3</sub>)  $\delta$ : 0.94 (3H, t, J=8), 1.65 (2H, m, J=8),  
2.38 (3H, s), 2.65 (2H, t, J=8), 7.22  
(1H, d, J=9), 7.98 (1H, dd, J=9, 2),  
8.15 (1H, s), 8.45 (1H, d, J=2).

-25-

Description 231,2-Dimethyl-5-nitro-3-propyl-1H-indole (D23)

5 The title compound was prepared in 89% yield from the nitroindole (D22) using a procedure similar to that in Description 2.

10 NMR (CDCl<sub>3</sub>)  $\delta$ : 0.90 (3H, t, J=8), 1.55 (2H, m, J=8),  
2.34 (3H, s), 2.70 (2H, t, J=8), 3.72  
(3H, s), 7.55 (1H, d, J=9), 7.92 (1H,  
dd, J=9, 2), 8.35 (1H, d, J=2).

Description 24

15

5-Amino-1,2-dimethyl-3-propyl-1H-indole (D24)

The title compound was prepared in 92% yield from the nitroindole (D23) using a procedure similar to that in  
20 Description 3.

NMR (CDCl<sub>3</sub>)  $\delta$ : 1.10 (3H, t, J=8), 1.75 (2H, m, J=8),  
2.42 (3H, s), 2.75 (2H, t, J=8), 3.65  
(3H, s), 3.95 (2H, s), 6.65 (1H, d,  
25 J=9), 6.92 (1H, s), 7.14 (1H, d, J=9).

Description 253-n-Hexyl-2-methyl-5-nitro-1H-indole (D25)

30

The title compound was prepared in 72% yield from the 4-nitrophenylhydrazones of 2-nonanone using the method of Fludzinski et al described in J.Med.Chem., 1986, 29, 2415.

-26-

NMR (CDCl<sub>3</sub>)  $\delta$ : 0.90 (3H, m), 1.30 (6H, m), 1.60 (2H, m), 2.42 (3H, s), 2.68 (2H, t, J=7), 7.22 (1H, m), 8.04 (1H, m), 8.20 (1H, s), 8.45 (1H, d, J=1).

5

Description 261,2-Dimethyl-3-n-hexyl-5-nitro-1H-indole (D26)

10 The title compound was prepared in 74% yield from the nitroindole (D25) using a procedure similar to that in Description 2.

15 NMR (CDCl<sub>3</sub>)  $\delta$ : 0.88 (3H, m), 1.30 (6H, m), 1.58 (2H, m), 2.35 (3H, s), 2.70 (2H, m), 3.65 (3H, s), 7.15 (1H, d, J=9), 7.94 (1H, m), 8.46 (1H, d, J=1).

Description 27

20

5-Amino-1,2-dimethyl-3-n-hexyl-1H-indole (D27)

The title compound was prepared in 84% yield from the nitroindole (D26) using a procedure similar to that in  
25 Description 3.

30 NMR (CDCl<sub>3</sub>)  $\delta$ : 0.88 (3H, m), 1.30 (6H, m), 1.55 (2H, m), 2.28 (3H, s), 2.62 (2H, t, J=8), 2.98 (2H, s), 3.55 (3H, s), 6.58 (1H, m), 6.80 (1H, d, J=1), 7.0 (1H, d, J=8).

-27-

Description 28Ethyl 2-Oxopentanoate (D28)

5 The sodium salt of 2-oxopentanoic acid (1.00g, 7.25 mM) was taken up in water and acidified to pH 1. The solution was extracted with ethyl acetate (3 x 100 ml), dried and solvents removed in vacuo. The resulting oil (0.67g) was taken up in ethanol (50 ml) and Amberlyst 15 added (0.67g).  
10 The suspension was stirred over 48h, the resin filtered off and solvents removed in vacuo to give the title compound as a slightly coloured oil (0.54g; 64%).

15 NMR (CDCl<sub>3</sub>)  $\delta$ : 0.95 (3H, t, J=7), 1.38 (3H, t, J=7), 1.55 (2H, m), 2.32 (2H, t, J=6), 4.32 (2H, q, J=7).

Description 2920 Ethyl 2-oxopentanoate 4-nitrophenylhydrazone (D29)

To a solution of the ester (D28) (0.53g, 3.6 mM) in ethanol (20 ml) was added 4-nitrophenylhydrazine (0.56g, 3.6 mM) and the suspension stirred for 0.5h. Concentrated hydrochloric  
25 acid (2 ml) was added to give a brown solution. After stirring for 0.5h the solution was cooled in ice and the precipitated title compound filtered off (0.72g; 69%).

30 NMR (CDCl<sub>3</sub>)  $\delta$ : 1.02 (3H, m), 1.40 (3H, m), 1.62 (2H, m), 2.60 (2H, m), 4.32 (2H, m), 7.28 (2H, m), 8.15 (2H, m), 8.30 (1H, s).

-28-

Description 303-Ethyl-5-nitro-1H-indole (D30)

5 The 4-nitrophenylhydrazone of ethyl 2-oxopentanoate (D29)  
(0.72g, 2.60 mM) was heated to reflux for 16h in  
concentrated hydrochloric acid. After cooling to room  
temperature the precipitated solid was filtered off.  
Chromatography on silica using dichloromethane as eluant  
10 gave the title compound as a yellow solid (0.22g; 45%).

NMR (CDCl<sub>3</sub>)  $\delta$ : 1.35 (3H, t, J=8), 2.80 (2H, q, J=7), 7.12  
(1H, m), 7.40 (1H, d, J=10), 8.12 (1H, dd,  
J=6, 1), 8.44 (1H, s), 8.60 (1H, m).

15

Description 313-Ethyl-1-methyl-5-nitro-1H-indole (D31)

20 The title compound was prepared in 95% yield from the  
corresponding indole (D30) following a procedure similar to  
that in Description 2.

NMR (CDCl<sub>3</sub>)  $\delta$ : 1.3 (3H, t, J=7), 2.0 (2H, q, J=7), 3.82  
25 (3H, s), 6.98 (1H, s), 7.27 (1H, d, J=8),  
8.12 (1H, dd, J=7, 1), 8.55 (1H, d, J=1).

Found: M<sup>+</sup> 204

C<sub>16</sub>H<sub>12</sub>N<sub>2</sub>O<sub>2</sub> requires 204.

30

-29-

Description 325-Amino-3-ethyl-1-methyl-1H-indole (D32)

5 The title compound was prepared from the corresponding nitro-indole (D31) in 98% yield following a procedure similar to that in Description 3.

10 NMR (CDCl<sub>3</sub>) δ: 1.25 (3H, m), 2.70 (2H, q, J=8), 3.64 (3H, s), 6.62 (1H, m), 6.71 (1H, s), 6.78 (1H, m), 7.06 (1H, m).

Description 3315 Phenyl N-(1-Methyl-1H-indol-5-yl)carbamate (D33)

To a solution of phenyl chloroformate (2.21 ml; 17.4 mM) in dry tetrahydrofuran (30 ml), cooled in a carbon tetrachloride / solid carbon dioxide bath, was added 5-amino-1-methylindole (D3) (2.31g; 15.8 mM) followed by triethylamine (2.40 ml; 17.4 mM). The mixture was stirred for 45 min at -20°C (bath temp.), then evaporated and the residue was dissolved in ethyl acetate, washed with brine, dried and evaporated to give the title compound (4.29g; 25 100%), m.p. 103-107°C (EtOAc/petrol).

NMR (CDCl<sub>3</sub>) δ: 3.80 (3H, s), 6.45 (1H, d, J=3), 6.93 (1H, broad s), 7.05 (1H, d, J=3), 7.25 (5H, m), 7.40 (2H, dd, J=8, 8), 7.74 (1H, broad s).

30

-30-

Description 342-Dimethylamino-5-nitropyridine (D34)

5 2-Chloro-5-nitropyridine (1.58g, 10 mM) was treated with a 33% w/w solution of dimethylamine in methylated spirit (18 ml, 100 mM). An exothermic reaction ensued, with formation of a yellow solid. After 0.5h the solid was filtered off. The filtrate was evaporated and the residue was combined  
10 with the yellow solid, and all material was dissolved in dichloromethane. This solution was washed with water and brine, dried and evaporated, to give the title compound (1.64g; 98%), m.p. 146-149°C.

15 NMR (CDCl<sub>3</sub>)  $\delta$ : 3.25 (6H, s), 6.48 (1H, d, J=10), 8.20 (1H, dd, J=10,3), 9.06 (1H, d, J=3).

Found: M<sup>+</sup> 167

C<sub>7</sub>H<sub>9</sub>N<sub>3</sub>O<sub>2</sub> requires 167.

20

Description 355-Amino-2-dimethylaminopyridine (D35)

25 2-Dimethylamino-5-nitropyridine (D34) (1.64g, 9.8 mM) was stirred with 10% palladium on charcoal (0.16g) in ethanol (200 ml) under 1 atmos. of hydrogen. After 6h the catalyst was filtered off onto Kieselguhr and the filtrate was evaporated. The residue was dissolved in diethyl ether,  
30 filtered again, and chromatographed on silica gel (50g) using ether as eluant. The eluted product was purified further by extraction with petrol (bp 60-80°C) to give the title compound as a reddish oil (0.66g; 49%).



-31-

NMR (CDCl<sub>3</sub>)  $\delta$ : 3.00 (6H, s), 6.47 (1H, d, J=9), 6.99 (1H, dd, J=9,3), 7.78 (1H, d, J=3).

Found: M<sup>+</sup> 137

5 C<sub>7</sub>H<sub>11</sub>N<sub>3</sub> requires 137

Description 36

3-Isopropyl-2-methyl-5-nitro-1H-indole (D36)

10

The title compound was prepared in 62% yield from the 4-nitrophenyl hydrazone of 4-methyl-2-pentanone using the method of Fludzinski *et al* described in J. Med. Chem., 1986, 29, 2415.

15

NMR (CDCl<sub>3</sub>)  $\delta$ : 1.42 (6H, d, J=6), 2.42 (3H, s), 3.20 (1H, m), 7.28 (1H, m), 8.03 (1H, dd, J=7,1), 8.12 (1H, s), 8.60 (1H, m).

20 Description 37

1,2-Dimethyl-3-isopropyl-5-nitro-1H-indole (D37)

The title compound was formed in 85% yield from the  
25 nitroindole (D36) following a procedure similar to that in Description 2.

NMR (CDCl<sub>3</sub>)  $\delta$ : 1.47 (6H, d, J=7), 2.42 (3H, s), 3.20 (1H, m), 3.68 (3H, s), 7.24 (1H, m), 8.04 (1H, m), 8.62 (1H, m).

30

Description 38

5-Amino-1,2-dimethyl-3-isopropyl-1H-indole (D38)

35 The title compound was formed in 57% yield from the nitroindole (D37) following a procedure similar to that in

-32-

## Description 3.

NMR (CDCl<sub>3</sub>)  $\delta$ : 1.35 (6H, d, J=7), 2.30 (3H, s), 3.15 (1H, m), 3.55 (3H, s), 6.60 (1H, m), 7.05 (2H, m).

5

Example 1N-(1,2-Dimethyl-3-ethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea hydrochloride (E1)

10

To a solution of the aminoindole (D1) (0.71g; 3.78mM) in dichloromethane (13.5ml) at 0°C was added a 12.5% solution of phosgene in toluene (3.28ml; 3.79mM). After stirring for 0.5h, triethylamine (1.15ml) was added and stirring was  
15 continued for 0.5h. A solution of 3-aminopyridine (0.34g; 3.6mM) in dichloromethane (10ml) was then added, and stirring continued for 3.5h at room temperature. Several drops of aqueous sodium hydroxide were added to the reaction mixture which was vigorously stirred for 0.5h. The reaction  
20 mixture was diluted with water and extracted with dichloromethane. The combined organic extracts were washed with brine, dried over sodium sulphate and evaporated to dryness. Chromatography on silica using dichloromethane as eluant afforded the title compound (0.2g; 17%) which was  
25 converted to the hydrochloride salt using hydrogen chloride in ether/ethanol, m.p. 158-165°C.

NMR (D<sub>6</sub>-DMSO)  $\delta$ :

1.12 (3H, t, J=8), 2.31 (3H, s), 2.63 (2H, q, J=8), 3.61  
30 (3H, s), 7.02 (1H, m), 7.28 (1H, d, J=10), 7.64 (1H, s),  
7.89 (1H, m), 8.31 (1H, m), 8.45 (1H, d, J=6), 9.13 (1H, s),  
9.22 (1H, s), 10.12 (1H, s).

Found: M<sup>+</sup> 308.1640

35 C<sub>18</sub>H<sub>20</sub>N<sub>4</sub>O requires 308.1637

-33-

Example 2N-(1-Methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea hydrochloride  
(E2)

5

Method A

The title compound was prepared from 5-amino-1-methyl-1H-indole(D3), phosgene and 3-aminopyridine using a  
10 procedure similar to that described for Example 1, in 27%  
yield m.p. 175-180°C.

NMR ( $d_6$ -DMSO)  $\delta$ : 3.76 (3H, s), 6.34 (1H, d, J=2), 7.16  
(1H, dd, J=8, 2), 7.29 (1H, d, J=2), 7.37 (1H, d, J=8),  
15 7.70 (1H, s), 7.87 (1H, dd, J=8, 8), 8.30 (1H, m), 8.45  
(1H, J=8), 9.08 (1H, m), 9.24 (1H, s), 10.03 (1H, s).

Found:  $M^+$  266.1667

$C_{15}H_{14}N_4O$  requires 266.1667

20

Method B

A solution of the aminoindole (D3) (1.95g; 13 mM) in  
25 dichloromethane (20 ml) was added dropwise to a solution of  
3-pyridyl isocyanate (D4) (prepared from 3-pyridinecarbonyl  
azide (2.14g; 15mM) in toluene) at room temperature. The  
reaction mixture was stirred for 17h, then cooled, and the  
precipitate filtered off to give the crude product (3.36g;  
30 95%). This was dissolved in hot ethanol and ethereal  
hydrogen chloride added to afford the title compound as its  
hydrochloride salt (3.1g; 80%) identical with the material  
prepared by method A.

-34-

Example 3N-(1,2,3-Trimethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea  
hydrochloride (E3)

5

The title compound was prepared in 51% yield from 5-amino-1,2,3-trimethyl-1H-indole (D6), phosgene and 3-aminopyridine using a procedure similar to that in Example 1, m.p. 330°C.

10 NMR (D<sub>6</sub>-DMSO) δ:

2.12 (3H, s), 2.30 (3H, s), 3.60 (3H, s), 7.04 (1H, dd, J=9, 2), 7.27 (1H, d, J=9), 7.58 (1H, d, J=2), 7.89 (1H, dd, J=9, 9), 8.32 (1H, m), 8.44 (1H, d, J=6), 9.11 15 (1H, d, J=2), 9.28 (1H, s), 10.22 (1H, s).

Found: M<sup>+</sup> 294.1485C<sub>17</sub>H<sub>18</sub>N<sub>4</sub>O requires: 294.148120 Example 4N-(1-Propyl-1H-indol-5-yl)-N'-(3-pyridyl)urea oxalate (E4)

The title compound was prepared in 43% yield from 5-amino-1-25 propyl-1H-indole (D8) and 3-pyridyl isocyanate (D4) using a procedure similar to that in Example 2 (Method B), the product being isolated as the oxalate salt, m.p. 165-169°C.

NMR (D<sub>6</sub> DMSO) δ:

30

0.82 (3H, t, J=7), 1.76 (2H, h, J=7), 4.10 (2H, t, J=7),

-35-

6.37 (1H, d, J=3), 7.13 (1H, d, J=9), 7.34 (1H, d, J=3),  
7.40 (1H, d, J=9), 7.44 (1H, m), 7.69 (1H, s), 8.05 (1H, m),  
8.24 (1H, d, J=6), 8.67 (1H, s), 8.73 (1H, d, J=2), 8.97  
(1H, s).

5

Found:  $M^+$  294.1485


$C_{17}H_{18}N_4O$  requires: 294.1481

### Example 5

10

N-(1-Methyl-1H-indol-4-yl)-N'-(3-pyridyl)urea hydrochloride  
(E5)

15



20

A solution of the aminoindole (D10) (0.44g; 3.01 mM) in dichloromethane (10 ml) was added dropwise to a solution of 3-pyridyl isocyanate (D4) (prepared from 3-pyridine carbonyl azide (0.51g; 3.4 mM) in toluene) at room temperature. The  
25 reaction mixture was stirred for 17h, then cooled and the precipitate filtered off to give the crude product (1g; 100%). This was dissolved in hot ethanol and ethereal hydrogen chloride added to afford the title compound as its hydrochloride salt (0.74g; 81%), m.p 238°C.

30

NMR ( $D_6$  DMSO)  $\delta$ :

3.79 (3H, s), 6.80 (1H, d, J=3), 7.11 (2H, dd, J=6, 6),  
7.30 (1H, d, J=3), 7.70 (1H, dd, J=6, 2), 7.90 (1H, m),  
35 8.33 (1H, d, J=6), 8.49 (1H, d, J=3), 9.13 (1H, d, J=2),  
9.40 (1H, s), 10.80 (1H, s).


-36-

Found:  $M^+$  266.1170 $C_{15}H_{14}N_4O$  requires: 266.1167Example 6

5

N-(1-Methyl-1H-indol-6-yl)-N'-(3-pyridyl)urea hydrochloride  
(E6)

10



E6

15

A solution of the aminoindole (D12) (0.3g; 2.05 mM) in dichloromethane (10 ml) was added dropwise to a solution of 3-pyridyl isocyanate (D4) (prepared from 3-pyridinecarbonyl azide (0.28g; 2.26 mM) in toluene) at room temperature. The reaction mixture was stirred for 17h, then cooled, and the precipitate filtered off to give the crude product (0.43g; 79%). This was dissolved in hot ethanol and ethereal hydrogen chloride added to afford the title compound as the hydrochloride salt (0.35g; 56%), m.p. 215°C.

25

NMR ( $CDCl_3$ )  $\delta$ :

3.73 (3H, s), 6.36 (1H, d,  $J=2$ ), 6.96 (1H, dd,  $J=11, 2$ ),  
7.24 (1H, d,  $J=2$ ), 7.45 (1H, d,  $J=11$ ), 7.75 (1H, d,  $J=3$ ),  
30 7.90 (1H, m), 8.32 (1H, d,  $J=8$ ), 8.46 (1H, d,  $J=3$ ), 9.13  
(1H, d,  $J=3$ ), 9.49 (1H, s), 10.25 (1H, s).

Found: C, 59.24; H, 4.96; N, 18.38.

 $C_{15}H_{15}N_4OCl$  requires: C, 59.50; H, 4.99; N, 18.51

35

-37-

Example 7N-(1H-Indol-5-yl)-N'-(3-pyridyl)urea hydrochloride (E7)5 Method A

A solution of commercially available 5-aminoindole (0.5g; 3.8 mM) in dichloromethane (5 ml) was added dropwise to a solution of 3-pyridyl isocyanate (D4) (prepared from 3-  
10 pyridinecarbonyl azide (0.62g; 4.2 mM) in toluene) at room temperature. The reaction mixture was stirred for 2 days, then cooled and the precipitate filtered off, to give the crude product (0.54g; 57%), which was dissolved in ethanol and converted to the hydrochloride salt using hydrogen  
15 chloride in ether, m.p. 180-185°C.

NMR (D<sub>6</sub>-DMSO)  $\delta$ : 6.38 (1H, s), 7.11 (1H, d, J=8), 7.35 (2H, m), 7.7 (1H, s), 7.92 (1H, m), 8.35 (1H, d, J=8), 8.49 (1H, d, J=3), 9.12 (1H, s), 9.39 (1H, s), 10.41 (1H, s), 11.7  
20 (1H, s).

Found: M<sup>+</sup> 252

C<sub>14</sub>H<sub>12</sub>N<sub>4</sub>O requires 252

25 Method B

Compound E7 may also be prepared by reacting 3-methyl-4-nitroaniline with 3-pyridyl isocyanate (D4) by the procedure of Method A. The resulting nitrophenyl urea may be  
30 subjected to a Leimgruber synthesis by condensation with dimethylformamide dimethylacetal with heating followed by hydrogenation over palladium and charcoal at high pressure to effect formation of the indole.

-38-

Example 8N-(1-Methyl-1H-indol-5-yl)-N'-methyl-N'-(3-pyridyl)urea  
(E8)

5

To a solution of carbonyl diimidazole (1.22g; 7.5 mM) in dichloromethane (10 ml) was added aminoindole (D3) (1.0g; 6.85 mM) in dichloromethane (10 ml). After stirring at room temperature for 15 min, the solution was evaporated to dryness. The residue was taken up in dimethylformamide (10 ml) and to this solution was added 3-methylaminopyridine (D13) (0.74g; 6.2 mM) in dimethylformamide (10 ml). The reaction mixture was heated to 90°C for 1h, then cooled and added dropwise to water (200 ml) with vigorous stirring. After cooling overnight, the precipitate was filtered and dried to give the crude product (1.99g). Chromatography on silica using dichloromethane as eluant afforded the title compound (0.81g; 42%), m.p. 58-60°C.

20 NMR (CDCl<sub>3</sub>) δ: 3.40 (3H, s), 3.75 (3H, s), 6.18 (1H, s), 6.39 (1H, d, J=3), 7.02 (1H, d, J=3), 7.09 (1H, dd, J=8, 1), 7.21 (1H, d, J=8), 7.42 (1H, m), 7.57 (1H, d, J=1), 7.75 (1H, m), 8.59 (1H, dd, J=3,1), 8.70 (1H, d, J=1).

25 Found: M<sup>+</sup> 280  
C<sub>16</sub>H<sub>16</sub>N<sub>4</sub>O requires 280

Example 930 N-Methyl-N-(1-methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea (E9)

The title compound was prepared from 1-methyl-5-methylamino-1H-indole (D15) and 3-pyridyl isocyanate (D4) using a procedure similar to that described for Example 2 Method B.

35 The crude product was obtained in 45% yield.



-39-

Recrystallisation from ethanol afforded the title compound, m.p. 168-170°C.

NMR (CDCl<sub>3</sub>)  $\delta$ : 3.39 (3H, s), 3.87 (3H, s), 6.35 (1H, s),  
5 6.55 (1H, d, J=3), 7.18 (3H, m), 7.43 (1H, d, J=8), 7.60  
(1H, d, J=1), 8.01 (1H, m), 8.2 (2H, m).

Found: C, 68.55; H, 5.79; N, 19.92%

C<sub>16</sub>H<sub>16</sub>N<sub>4</sub>O requires C, 68.55; H, 5.75; N, 19.99%

10

Example 10

N-Methyl-N-(1-methyl-1H-indol-5-yl)-N'-methyl-N'-(3-pyridyl)urea (E10)

15


To a suspension of 80% sodium hydride (0.06g; 2 mM) in dimethylformamide (5 ml), was added the monomethyl urea (E9) (0.5g; 1.79 mM). After stirring at room temperature for 0.5h, methyl iodide (0.12 ml; 1.93 mM) was added dropwise.  
20 Stirring was continued at room temperature for 1h, then heated at 50°C for 1h. The reaction mixture was cooled in ice, then quenched with water. The mixture was then extracted with dichloromethane, washed with water, dried over sodium sulphate and evaporated to give the crude  
25 product (0.59g). Chromatography on silica using dichloromethane as eluant afforded the title compound (0.31g; 60%) which was recrystallised from cyclohexane to give a white solid (160 mg) m.p. 91-92.5°C.

30 NMR (CDCl<sub>3</sub>)  $\delta$ : 3.18 (3H, s), 3.25 (3H, s), 3.71 (3H, s), 6.3  
(1H, d, J=3), 6.6 (1H, dd, J=8,1), 6.85 (1H, m), 7.01 (4H, m), 8.10 (2H, m).

Found: C, 69.57; H, 6.21; N, 19.04%

35 C<sub>17</sub>H<sub>18</sub>N<sub>4</sub>O requires C, 69.37; H, 6.16; N, 19.03%

-40-

Example 11N-(1-Methyl-1H-indol-5-yl)-N'-(2-pyridyl)urea (E11)

10 The title compound was prepared from 5-amino-1-methylindole (D3) and 2-aminopyridine using a procedure similar to that described for Example 8. The crude product was obtained in 83% yield. Recrystallisation from ethanol afforded the title compound in 70% yield, m.p. 182-185°C.

15

NMR (CDCl<sub>3</sub>)  $\delta$ : 3.8 (3H, s), 6.42 (1H, d, J=3), 6.9 (1H, m), 7.05 (1H, d, J=1), 7.2 (1H, d, J=8), 7.25 (1H, d, J=1), 7.32 (1H, dd, J=8, 1), 7.61 (1H, m), 7.88 (1H, s), 8.25 (1H, d, J=3), 9.11 (1H, s), 11.18 (1H, s).

20

Found: M<sup>+</sup> 266

C<sub>15</sub>H<sub>14</sub>N<sub>4</sub>O requires 266.

Example 12

25

N-(1,4-Dimethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea hydrochloride (E12)

The title compound was prepared from 5-amino-1,4-dimethylindole (D17) and 3-pyridyl isocyanate (D4) following the procedure described in Example 2 Method B. This gave a yellow-green powder in 21% yield.

35 NMR (D<sub>6</sub>-DMSO)  $\delta$ : 2.38 (3H, s), 3.76 (3H, s), 6.45 (1H, d, J=3), 7.24 (2H, s), 7.30 (1H, d, J=3), 7.89 (1H, dd, J=8, 5), 8.33 (1H, d, J=8), 8.44 (1H, d, J=5), 8.67 (1H, s), 9.11 (1H, fine d), 10.3 (1H, b s).

-41-

Found:  $M^+$  280 $C_{16}H_{16}N_4O$  requires 280

Found: C, 57.8; H, 5.5; N, 16.9%.  $C_{16}H_{16}N_4O \cdot HCl \cdot H_2O$   
5 requires C, 54.4, H, 5.7; N, 16.7%

Example 13

N-(1-Methyl-1H-indol-5-yl)-N'-(2-chloropyrid-3-yl)urea  
10 hydrochloride (E13)

A stirred suspension of carbonyl diimidazole (0.34g, 2.1 mM) in dry dichloromethane (5 ml) was treated with a solution of 5-amino-1-methyl-1H-indole (D3) (0.29g, 2 mM) in dry  
15 dichloromethane (5 ml). After 0.25h the reaction mixture was evaporated to dryness, and the residue dissolved in dimethylformamide (10 ml). 3-Amino-2-chloro-pyridine (0.23g, 22 mM) was added to the reaction mixture which was heated to 90°C for 1h, then cooled and added to water (200  
20 ml) with vigorous stirring. The precipitate was filtered, dried and recrystallised from ethanol affording the title compound as an off white solid (0.25g; 42%) which was converted to the hydrochloride salt using hydrogen chloride in ether, m.p. 155°C.

25

NMR ( $D_6$ -DMSO)  $\delta$ : 3.78 (3H, s), 6.37 (1H, d, J=5), 7.15 (1H, dd, J=12, 3), 7.30 (1H, d, J=5), 7.40 (2H, m), 7.72 (1H, d, J=3), 8.02 (1H, d, J=5), 8.49 (1H, d, J=3), 8.6 (1H, d, J=12), 9.34 (1H, s).

30

Found:  $M^+$  299, 301 $C_{15}H_{13}N_4O$  Cl requires 299, 301

-42-

Example 14N-(1-Methyl-1H-indol-5-yl)-N'-(2-chloropyrid-5-yl)urea hydrochloride (E14)

5


The title compound was prepared in 60% yield from 5-amino-1-methyl-1H-indole (D3), carbonyl diimidazole and 5-amino-2-chloropyridine, using a procedure similar to that described in Example 13. m.p. 212°C.

10

NMR ( $D_6$ -DMSO)  $\delta$ : 3.78 (3H, s), 6.32 (1H, d, J=5), 7.15 (1H, dd, J=12,3), 7.28 (1H, d, J=5), 7.40 (2H, m), 7.70 (1H, d, J=3), 8.00 (1H, dd, J=12,5), 8.50 (1H, d, J=5).

15 Found:  $M^+$  299, 301 $C_{15}H_{13}N_4O$  Cl requires 299, 301.Example 1520 N-(1-Methyl-1H-indol-5-yl)-N'-(3-hydroxypyrid-2-yl)urea hydrochloride (E15)

25



N-(1-Methyl-1H-indol-5-yl)-N'-(3-benzyloxy-pyrid-2-yl)urea (D18) (0.37g, 1 mM) was hydrogenated for 2h in ethanol (40 ml) at atmospheric pressure and room temperature. The reaction mixture was filtered through kieselguhr, washed with ethanol. The filtrate was evaporated in vacuo to afford the title compound (0.21g, 74%) which was converted to the hydrochloride salt using hydrogen chloride in ether. 35 m.p. 223°C.

-43-

NMR ( $D_6$ -DMSO)  $\delta$ : 3.78 (3H, s), 6.39 (1H, d, J=5), 6.95 (1H, m), 7.23 (2H, m), 7.3 (1H, d, J=5), 7.39 (1H, d, J=12), 7.83 (2H, m), 7.95 (1H, s).

5 Found:  $M^+$  282

$C_{15}H_{14}N_4O_2$  requires 282

Example 16

10 N-(1,3-Diethyl-2-methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea  
(E16)

The title compound was prepared from the aminoindole (D21) and 3-pyridyl isocyanate (D4) using a procedure similar to  
15 that described for Example 2 Method B. The crude product was obtained in 79% yield. Recrystallisation from ethanol afforded the title compound, m.p. 192-193°C.

NMR ( $D_6$  DMSO)  $\delta$ : 1.12 (3H, t, J=8), 1.20 (3H, t, J=8), 2.31  
20 (3H, s), 2.65 (2H, q, J=8), 4.09 (2H, q, J=8), 7.02 (1H, dd, J=9,3), 7.26 (1H, d, J=9), 7.29 (1H, m), 7.63 (1H, d, J=3), 7.98 (1H, m), 8.17 (1H, m), 8.55 (1H, s), 8.60 (1H, d, J=3), 8.74 (1H, s).

25 Example 17

N-(1,2-Dimethyl-3-propyl-1H-indol-5-yl)-N'-(3-pyridyl)urea  
hydrochloride (E17)

30 The title compound was prepared from the aminoindole (D24) and 3-pyridyl isocyanate (D4) using a procedure similar to that described for Example 2 Method B, m.p. 132-134°C.

NMR ( $D_6$ -DMSO)  $\delta$ : 0.88 (3H, t, J=8), 1.54 (2H, m), 2.28 (3H,  
35 s), 2.58 (2H, m), 3.62 (3H, s), 7.04 (1H, d, J=4), 7.28 (1H,

-44-

d, J=6), 7.60 (1H, s), 7.90 (1H, m), 8.32 (1H, d, J=4), 8.45 (1H, d, J=6), 9.12 (1H, s), 9.25 (1H, s), 10.22 (1H, s).

Found:  $M^+$  322

5  $C_{19}H_{22}N_4O$  requires 322

### Example 18

N-(1,2-Dimethyl-3-n-hexyl-1H-indol-5-yl)-N'-(3-pyridyl)urea


10 (E18)

The title compound was prepared from the aminoindole (D27) and 3-pyridyl isocyanate (D4) using a procedure similar to that described for Example 2 Method B, the product being  
15 isolated as the free base.

NMR ( $D_6$  DMSO)  $\delta$ : 0.84 (3H, m), 1.25 (6H, m), 1.52 (2H, m), 2.30 (3H, s), 2.62 (2H, m), 3.58 (3H, s), 7.05 (1H, dd, J=8, 2), 7.28 (1H, d, J=6), 7.60 (1H, s), 7.90 (1H, m), 8.32 (1H, 20 m), 8.45 (1H, d, J=6), 9.15 (1H, m), 9.25 (1H, s), 10.15 (1H, s).

### Example 19

25 N-(1-Methyl-1H-indol-4-yl)-N'-(4-pyridyl)urea hydrochloride  
(E19)



30

4-Amino-1-methyl-1H-indole (D10) (0.44g) was treated successively with phosgene (solution in toluene) and 4-aminopyridine as described in Example 1. The reaction  
35

-45-

mixture was partitioned between dichloromethane and water, and filtered. The solid, the crude free base of the title compound, was filtered off and dried in vacuo. The filtrate was separated, and the organic portion was washed with  
5 brine, dried and evaporated to give an oil. The oil was chromatographed on silica using methanol/chloroform (0-10% methanol, gradient) as eluant, giving further crude free base.

10 The two portions of free base were combined, and this material (0.49g) was suspended in ethanol (50 ml) at reflux. Briefly after removing from the steam bath, HCl in ether (1.1 M, 3 ml) was added. The suspension was brought back to reflux, and then cooled. Filtration and drying gave the  
15 title compound (0.30g) as a grey-brown solid.

NMR ( $D_6$ -DMSO)  $\delta$ : 3.80 (3H, s), 6.87 (1H, d, J=3), 7.15 (2H, m), 7.32 (1H, d, J=3), 7.73 (1H, d, J=7), 7.95 (2H, d, J=6), 8.60 (2H, d, J=6), 9.76 (1H, s), 11.84 (1H, s), 14.5 (v  
20 broad).

Found:  $M^+$  266,  $C_{15}H_{14}N_4O$  requires 266

Found: C, 55.89; H, 5.15; N, 17.20%  
25  $C_{15}H_{14}N_4O \cdot HCl \cdot H_2O$  requires C, 56.16; H, 5.34; N, 17.47%

#### Example 20

N-(3-Ethyl-1-methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea (E20)

30

The title compound was prepared in 82% yield from the aminoindole (D32) and 3-pyridyl isocyanate (D4) using a procedure similar to that described for Example 2, Method B, the product being isolated as the free base.

35

-46-


NMR (CDCl<sub>3</sub>)  $\delta$ : 1.26 (3H, t, J=7), 2.68 (2H, q, J=7), 3.69 (3H, s), 6.85 (1H, s), 7.04 (1H, m), 7.30 (2H, m), 7.56 (1H, s), 7.95 (1H, m), 8.15 (1H, m), 8.65 (2H, m), 8.80 (1H, s).

5 Found: M<sup>+</sup> 294, C<sub>17</sub>H<sub>18</sub>N<sub>4</sub>O requires 294

Example 21

N-(1,2-Dimethyl-3-ethyl-1H-indol-5-yl)-N'-(2-pyridyl)urea

10 (E21)



The title compound was prepared from the aminoindole (D1),  
20 phosgene and 2-aminopyridine using a procedure similar to  
that described for Example 1, the product being isolated as  
the free base, m.p. 120-123°C.

NMR (CDCl<sub>3</sub>)  $\delta$ : 1.22 (3H, t, J=8), 2.35 (3H, s), 2.73  
25 (2H, q, J=8), 3.64 (3H, s), 6.91 (2H, m), 7.19 (1H, d, J=9),  
7.28 (1H, m), 7.60 (1H, m), 7.77 (1H, m), 8.27 (2H, m), 11.5  
(1H, broad s).

Found: C, 70.03; H, 6.36; N, 17.97%  
30 C<sub>18</sub>H<sub>20</sub>N<sub>4</sub>O requires C, 70.11; H, 6.54; N, 18.17%




-47-

Example 22

N-(1,2-Dimethyl-3-ethyl-1H-indol-5-yl)-N'-(4-pyridyl)urea  
hydrochloride (E22)

5



10

(E22)

The title compound was prepared in 47% yield from the aminoindole (D1), phosgene and 4-aminopyridine using a procedure similar to that described for Example 1, the product being isolated as the hydrochloride salt, m.p. 237-243°C.

NMR (D<sub>6</sub> DMSO)  $\delta$ : 1.12 (3H, t, J=8), 2.30 (3H, s), 2.62 (2H, q, J=8), 3.61 (3H, s), 7.05 (1H, dd, J=9,2), 7.29 (1H, d, J=9), 7.64 (1H, d, J=2), 7.90 (2H, d, J=6), 8.57 (2H, d, J=6), 9.67 (1H, broad s), 11.28 (1H, broad s).

Found: M<sup>+</sup> 308  
C<sub>18</sub>H<sub>20</sub>N<sub>4</sub>O requires 308

25

Example 23

N-(1-Methyl-1H-indol-5-yl)-N'-(2-dimethylamino-5-pyridyl)urea (E23)

30

5-Amino-2-dimethylaminopyridine (D35) (0.137g; 1 mM) was stirred with 80% sodium hydride (66 mg; 2.2 mM) in dry dimethylformamide (5 ml) for 15 min at room temperature under nitrogen. The phenyl carbamate (D33) was then added and the mixture was stirred overnight at room temperature.

35

-48-

Solvent was then removed in vacuo and the residue was dissolved in dichloromethane/methanol, washed with water and brine, dried and evaporated. The residue was triturated with dichloromethane/petrol, and the solid material was chromatographed on silica gel and eluted with 2% methanol/dichloromethane. This gave the title compound (60 mg; 19%), m.p. 220-226°C.

NMR ( $D_6$  DMSO)  $\delta$ : 2.98 (6H, s), 3.75 (3H, s), 6.32 (1H, d, J=3), 6.62 (1H, d, J=9), 7.11 (1H, d, J=8), 7.26 (1H, d, J=3), 7.31 (1H, d, J=8), 7.66 (2H, m), 8.10 (1H, d, J=3), 8.20 (1H, s), 8.38 (1H, s).

Found:  $M^+$  309

$C_{17}H_{19}N_5O$  requires 309

#### Example 24

N-(1,2-Dimethyl-3-isopropyl-1H-indol-5-yl)-N'-(3-pyridyl)urea hydrochloride (E24)

The title compound was prepared from the aminoindole (D38) and 3-pyridyl isocyanate (D4) using a procedure similar to that described for Example 2, Method B.

NMR ( $D_6$  DMSO)  $\delta$ : 1.42 (6H, d, J=7), 2.42 (3H, s), 3.22 (1H, m), 3.68 (3H, s), 7.12 (1H, m), 7.36 (1H, m), 7.90 (1H, s), 7.98 (1H, m), 8.38 (1H, m), 8.55 (1H, m), 9.20 (1H, s), 9.30 (1H, s), 10.22 (1H, s).

Found  $M^+$  322

$C_{19}H_{22}N_4O$  requires 322.

-49-

Example 25N-(1,3-Diethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea (E25)

5 The title compound was prepared from ethyl 1,3-diethyl-5-nitro-1H-indole-2-carboxylate by hydrolysis and decarboxylation, then using a procedure similar to that in Description 3 and Example 2, Method B, the product being isolated as the free base.

10

m.p. 164-165°C

NMR (CDCl<sub>3</sub>) δ: 1.28 (3H, t, J=7), 1.45 (3H, t, J=7), 2.74 (2H, q, J=7), 4.12 (2H, q, J=7), 6.82 (1H, bs), 6.95 (1H, s), 7.10 (2H, m), 7.25 (2H, m), 7.58 (1H, s), 8.07 (1H, m), 8.24 (1H, m), 8.32 (1H, m).

Found: M<sup>+</sup> 308

C<sub>18</sub>H<sub>20</sub>N<sub>4</sub>O requires 308

20

Found: C, 69.93; H, 6.38; N, 17.98%

C<sub>18</sub>H<sub>20</sub>N<sub>4</sub>O requires C, 70.11; H, 6.54; N, 18.17%

Example 26

25

N-(3-Isopropyl-1-methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea (E26)

The title compound was prepared from ethyl 3-isopropyl-1-methyl-5-nitro-1H-indole-2-carboxylate by hydrolysis and decarboxylation, then using a procedure similar to that in Description 3 and Example 2, Method B, the product being isolated as the free base.

35 NMR (CDCl<sub>3</sub>) δ: 1.32 (6H, d, J=6), 3.15 (1H, m), 3.76 (3H, s), 6.76 (1H, bs), 6.88 (1H, s), 7.02 (1H, m), 7.13 (1H, m),

-50-

7.25 (2H, m), 7.60 (1H, m), 8.08 (1H, d, J=8), 8.22 (1H, m),  
8.30 (1H, m).

Found:  $M^+$  308

5  $C_{18}H_{20}N_4O$  requires 308

### Example 27

#### N-(1,3-Dimethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea (E27)

10

The title compound was prepared from ethyl 1,3-dimethyl-5-nitro-1H-indole-2-carboxylate by hydrolysis and decarboxylation, then using a procedure similar to that in Description 3 and Example 2, Method B, the product being  
15 isolated as the free base.

m.p. 210°C

NMR ( $D_6$  - DMSO)  $\delta$ : 2.25 (3H, s), 3.72 (3H, s), 6.88 (1H,  
20 s), 7.10 (1H, dd, J=9, 1), 7.22 (2H, m), 7.69 (1H, d, J=1),  
8.04 (1H, m), 8.16 (1H, m), 8.31 (1H, s), 8.68 (1H, m).

Found:  $M^+$  280

$C_{16}H_{16}N_4O$  requires 280

25

TABLE 1


								Salt
	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	R <sub>7</sub>	
E1	CH <sub>3</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	H	H	H	H	HCl
E2	CH <sub>3</sub>	H	H	H	H	H	H	HCl
E3	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	H	H	H	H	HCl
E4	(CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>	H	H	H	H	H	H	(COOH) <sub>2</sub>
E7	H	H	H	H	H	H	H	HCl
E8	CH <sub>3</sub>	H	H	H	CH <sub>3</sub>	H	H	-
E9	CH <sub>3</sub>	H	H	H	H	CH <sub>3</sub>	H	-
E10	CH <sub>3</sub>	H	H	H	CH <sub>3</sub>	CH <sub>3</sub>	H	-
E12	CH <sub>3</sub>	H	H	H	H	H	4-CH <sub>3</sub>	HCl
E13	CH <sub>3</sub>	H	H	2-Cl	H	H	H	HCl
E14	CH <sub>3</sub>	H	H	6-Cl	H	H	H	HCl
E16	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	H	H	H	H	-
E17	CH <sub>3</sub>	CH <sub>3</sub>	nC <sub>3</sub> H <sub>7</sub>	H	H	H	H	HCl
E18	CH <sub>3</sub>	CH <sub>3</sub>	nC <sub>6</sub> H <sub>13</sub>	H	H	H	H	-

TABLE 1 (CONTINUED)

	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	R <sub>7</sub>	Salt
E20	CH <sub>3</sub>	H	C <sub>2</sub> H <sub>5</sub>	H	H	H	H	-
E23	CH <sub>3</sub>	H	H	6-N(CH <sub>3</sub> ) <sub>2</sub>	H	H	H	-
E24	CH <sub>3</sub>	CH <sub>3</sub>	iC <sub>3</sub> H <sub>7</sub>	H	H	H	H	HCl
E25	C <sub>2</sub> H <sub>5</sub>	H	C <sub>2</sub> H <sub>5</sub>	H	H	H	H	-
E26	CH <sub>3</sub>	H	iC <sub>3</sub> H <sub>7</sub>	H	H	H	H	-
E27	CH <sub>3</sub>	H	CH <sub>3</sub>	H	H	H	H	-

-53-

Pharmacological data[<sup>3</sup>H]-mesulergine binding to pig choroid plexus membranes  
in vitro

5

Evidence from the literature suggests that 5-HT<sub>1C</sub> antagonists may have a number of therapeutic indications including the treatment of anxiety, migraine, depression, feeding disorders and obsessive compulsion disorders.

- 10 (Curzon and Kennett, 1990; Fozard and Gray, 1989) and Alzheimer's Disease (Lawlor, 1989, J. Arch. Gen. Psychiat. Vol. 46 p.542).

The affinity of test drugs for the 5-HT<sub>1C</sub> binding site can  
15 be determined by assessing their ability to displace [<sup>3</sup>H]-mesulergine from 5-HT<sub>1C</sub> binding sites in pig choroid plexus membranes. The method employed was similar to that of Pazos et al, 1984.

- 20 Pooled pig choroid plexi were homogenised in 20 vols of Tris HCl buffer (pH7.4) (containing 4mM CaCl<sub>2</sub> and 0.01% ascorbic acid) and centrifuged at 50,000g for 15 min at 4°C. The supernatant was removed and re-centrifuged. This was repeated a further two times with the incubation of the  
25 homogenate (37°C for 15 min) before the final centrifugation. The final pellet was resuspended in 20vols of buffer and stored at -70°C until use.

- The tissue suspension (50µl) was incubated with  
30 [<sup>3</sup>H]-mesulergine (2nM) in Tris HCl buffer (pH7.4) at 37°C (containing 0.01% ascorbic acid, 4mM CaCl<sub>2</sub>) and 3 x 10<sup>-8</sup>M spiperone for 30 minutes. Non-specific binding was measured in the presence of mianserin (10<sup>-6</sup>M). Six concentrations of test drug (10<sup>-9</sup> to 10<sup>-4</sup>M final concentration) were added in  
35 a volume of 50µl. The total assay volume was 500µl. Incubation was stopped by rapid filtration using a Skatron

-54-

cell harvester and radioactivity measured by liquid scintillation spectrometry. The  $IC_{50}$  values were determined and the  $pK_i$  (the negative logarithm of the inhibition constant) calculated from the Cheng Prusoff equation where

$$K_i = \frac{IC_{50}}{1 + \frac{C}{K_d}}$$

$K_i$  = inhibition constant.

$C$  = concentration of [ $^3H$ ]-mesulergine

$K_d$  = Affinity of mesulergine for 5-HT $_1C$  binding sites.

Curzon, G.A. and Kennett, G.A. (1990). TIPS, Vol. 11, 181-182.

Fozard, J.R. and Gray, J.A. (1989). TIPS, Vol. 10, 307-309.

Pazos, A. et al. (1984). Eur. J. Pharmacol., 106, 531-538.

Results are shown in Table 2.

## Table 2

	<u>Compound</u>	[ $^3H$ ]-Mesulergine <u><math>pK_i</math></u>
25	E1	7.6
	E2	6.8
	E3	6.7
	E4	6.7
	E5	6.7
30	E6	6.5

Compounds of the remaining examples have a  $pK_i > 5$ .



-55-

Rat stomach fundus

5-Hydroxytryptamine (5-HT) induces contractions of the rat stomach fundus through a 5-HT receptor that has the characteristics of a 5-HT<sub>1C</sub> receptor (Blackburn et al, 1990). Hence, this tissue can be used to assess the 5-HT<sub>1C</sub> antagonist actions of test drugs.

Rat stomach strips (6 x 4mm) were suspended under a 4g tension in 5ml baths containing Tyrode solution, gassed with a mixture of 95% O<sub>2</sub>/5% CO<sub>2</sub>. After a 1 hour equilibration period, two dose response curves were constructed to 5-HT (final concentrations, 10<sup>-9</sup> to 3 x 10<sup>-6</sup>M). Test drugs were then incubated at a final concentration of 10<sup>-6</sup>M for 30 mins and another dose-response curve constructed to 5-HT. The apparent dissociation constant of a test drug, K<sub>B</sub>, can be calculated from the equation where  $K_B = \frac{[B]}{DR-1}$

where B = concentration of the test drug and DR = the dose ratio (the factor by which the concentration of the agonist has to be increased in the presence of the test drug to obtain an identical effect observed in the absence of the test drug).

The results are shown in Table 3

Blackburn et al. (1990). Eur. J. Pharmacol., 180, 229-237.

Table 3

	<u>Compound</u>	<u>K<sub>B</sub></u>
5	E1	$3.2 \times 10^{-8}M$
	E2	$1 \times 10^{-7}M$
	E3	$2.5 \times 10^{-8}M$
	E4	$4.0 \times 10^{-7}M$
	E5	$2.5 \times 10^{-7}M$
10	E6	$1.7 \times 10^{-7}M$

Social Interaction Test

Potential anxiolytic properties have been evaluated using the social interaction test based on that described by File (1980 J.Neurosci.Meth., 2, 219). Active social interaction between male rats is usually quantitated by counting interactive behaviours such as following, grooming, sniffing, climbing over or under, biting, mounting and boxing. This behaviour is suppressed when the rats encounter each other in an environment which is novel and brightly lit. Under these circumstances anxiolytic drugs will enhance the level of social interaction.

Rats were housed in groups of 8 in a holding room adjacent to the experimental chamber for 8 days. They were then housed singly in the same room for 3 days prior to the experimental day. On the experimental day rats were injected p.o. 1h pretest with vehicle or drug in pairs at 15 min intervals beginning at 10.00 am. 60 Mins later they were placed with a weight matched pair mate (encountered for the first time) in the social interaction box in a separate room. The box was made of white perspex 54 x 37 x 26 cm

-57-

with no lid. The floor was divided into 24 equal squares and the cage was brightly lit. Active social interaction was scored blind over the next 15 min by remote video monitoring to give total interaction scores. The number of squares crossed by each rat was also scored and summed. At the end of each test the box was carefully wiped with a damp cloth. Unlike anxiolytic drugs, treatments that enhance social interaction by stimulant action will also increase locomotion. Treatments that are sedative reduce locomotion.

10

### Test Results

The compound of Example 2 showed a significant increase in social interaction at doses of 2-40 mg/kg.

15

### Geller-Seifter Procedure

Potential anxiolytic properties are evaluated using the Geller-Seifter procedure based on that originally described by Geller and Seifter, (1960) Psychopharmacologia, 1, 482-492. This procedure has been shown to be selective for drugs with anxiolytic properties (Cook and Sepinwall, (1975) 'Mechanism of Action of Benzodiazepines' ed. Costa, E. and Greengard, P., Raven Press, New York, pp. 1-28).

25

Rats are trained on a variable interval 30 sec schedule (VI30) to press a lever in order to obtain food reward. The 5 min sessions of the VI30 schedule alternate with 2-5 min of a schedule (FR5) in which every 5th lever press is followed by presentation of a food pellet paired with a 0.5 sec mild footshock. The total study lasts approximately 30 mins. Rats typically respond with high rates of lever pressing under the VI30 schedule and low response rates under the FR5 'conflict' session. Anxiolytic drugs increase

-58-

the suppressed response rates of rats in a 'conflict' session.

Drugs are administered intraperitoneally or orally to groups of 3-8 rats 30 min before testing.


The results are expressed as the percentage increase in the square root of the total number of lever presses in the FR5 'conflict' session. Square root transformation is necessary to normalise the data for statistical analysis using parametric methods.

The compound of Example 2 showed a significant increase in responding in the 'conflict' session at dose levels in the range 5-40 mg/kg p.o.

Claims

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:

5



15

wherein:

R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are independently hydrogen or C<sub>1-6</sub> alkyl;  
 R<sub>4</sub> is hydrogen, C<sub>1-6</sub> alkyl, halogen, hydroxy or NR<sub>8</sub>R<sub>9</sub> where  
 20 R<sub>8</sub> and R<sub>9</sub> are independently hydrogen or C<sub>1-6</sub> alkyl;  
 R<sub>5</sub> and R<sub>6</sub> are independently hydrogen or C<sub>1-6</sub> alkyl; and  
 R<sub>7</sub> is hydrogen, C<sub>1-6</sub> alkyl or halogen; and wherein the urea  
 moiety is attached at the 4-, 5- or 6-position of the indole  
 ring.

25

2. A compound according to claim 1 wherein the urea  
 moiety is attached at the 3-, 4- or 5-position of the  
 pyridine ring.

30 3. A compound according to claim 1 or 2 wherein the urea  
 moiety is attached at the 4- or 5-position of the indole  
 ring.

4. A compound according to any preceding claim wherein  
 35 any alkyl moiety within variables R<sub>1</sub> to R<sub>9</sub> is C<sub>1-3</sub> alkyl.

-60-

5. A compound according to claim 4 wherein  $R_1$  is methyl,  $R_2$  is methyl or hydrogen,  $R_3$  is hydrogen, methyl, ethyl, n-propyl or iso-propyl,  $R_4$  is hydrogen and  $R_5$ ,  $R_6$  and  $R_7$  are independently hydrogen or methyl.
- 5 6. N-(1,2-Dimethyl-3-ethyl-1H-indol-5-yl)-N'-(3-pyridyl)-urea.
7. N-(1-Methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
- 10 8. N-(1,2,3-Trimethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
9. N-(1-Propyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
- 15 10. N-(1-Methyl-1H-indol-4-yl)-N'-(3-pyridyl)urea.
11. N-(1-Methyl-1H-indol-6-yl)-N'-(3-pyridyl)urea.
12. N-(1H-Indol-5-yl)-N'-(3-pyridyl)urea.
- 20 13. N-(1-Methyl-1H-indol-5-yl)-N'-methyl-N'-(3-pyridyl)urea.
14. N-Methyl-N-(1-methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
- 25 15. N-Methyl-N-(1-methyl-1H-indol-5-yl)-N'-methyl-N'-(3-pyridyl)urea.
- 30 16. N-(1-Methyl-1H-indol-5-yl)-N'-(2-pyridyl)urea.
17. N-(1,4-Dimethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
18. N-(1-Methyl-1H-indol-5-yl)-N'-(2-chloropyrid-3-yl)urea.
- 35

-61-

19. N-(1-Methyl-1H-indol-5-yl)-N'-(2-chloropyrid-5-yl)urea.
20. N-(1-Methyl-1H-indol-5-yl)-N'-(3-hydroxypyrid-2-5 yl)urea.
21. N-(1,3-Diethyl-2-methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
- 10 22. N-(1,2-Dimethyl-3-propyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
23. N-(1,2-Dimethyl-3-n-hexyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
- 15 24. N-(1-Methyl-1H-indol-4-yl)-N'-(4-pyridyl)urea.
25. N-(3-Ethyl-1-methyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
- 20 26. N-(1,2-Dimethyl-3-ethyl-1H-indol-5-yl)-N'-(2-pyridyl)urea.
27. N-(1,2-Dimethyl-3-ethyl-1H-indol-5-yl)-N'-(4-pyridyl)urea.
- 25 28. N-(1-Methyl-1H-indol-5-yl)-N'-(2-dimethylamino-5-pyridyl)urea.
29. N-(1,2-Dimethyl-3-isopropyl-1H-indol-5-yl)-N'-(3-30 pyridyl)urea.
30. N-(1,3-Diethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.
31. N-(3-Isopropyl-1-methyl-1H-indol-5-yl)-N'-(3-35 pyridyl)urea.

-62-

32. N-(1,3-Dimethyl-1H-indol-5-yl)-N'-(3-pyridyl)urea.


33. A pharmaceutically acceptable salt of a compound according to any one of claims 6 to 32.

5

34. A process for the preparation of a compound according to claim 1, which process comprises:

(a) the coupling of a compound of formula (II);

10




15

(II)

with a compound of formula (III);

20



25

(III)

wherein B is attached at the 4-, 5- or 6-position of the indole ring and A and B contain the appropriate functional group(s) necessary to form the moiety -NR<sub>5</sub>'CONR<sub>6</sub>'- when coupled, wherein R<sub>5</sub>' and R<sub>6</sub>' are R<sub>5</sub> and R<sub>6</sub> as defined in claim 1 or groups convertible thereto, and the variables R<sub>1</sub>', R<sub>2</sub>', R<sub>3</sub>', R<sub>4</sub>' and R<sub>7</sub>' are R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub> and R<sub>7</sub> respectively, as defined in claim 1, or groups convertible thereto, and thereafter optionally and as necessary and in

30


35



-63-

any appropriate order, converting any  $R_1'$ ,  $R_2'$ ,  $R_3'$ ,  $R_4'$ ,  $R_5'$ ,  $R_6'$  and  $R_7'$  when other than  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$  respectively to  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$ , interconverting  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$ , and forming a pharmaceutically acceptable salt thereof, or

(b) cyclising a compound of formula (IV):




15

wherein  $R_4'$ ,  $R_5'$ ,  $R_6'$  and  $R_7'$  are as defined in formulae (II) and (III) and C and D contain the appropriate functional group(s) necessary to form the indole ring substituted by  $R_1'$ ,  $R_2'$  and  $R_3'$  as defined in formula (III), and thereafter optionally and as necessary in any appropriate order, converting any  $R_1'$ ,  $R_2'$ ,  $R_3'$ ,  $R_4'$ ,  $R_5'$ ,  $R_6'$  and  $R_7'$  when other than  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$ , to  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$ , interconverting  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$  and forming a pharmaceutically acceptable salt.

20

25

35. A compound of formula (IV):



-64-

wherein  $R_4'$ ,  $R_5'$ ,  $R_6'$  and  $R_7'$  are  $R_4$ ,  $R_5$ ,  $R_6$  and  $R_7$  as defined in claim 1 or groups convertible thereto and C and D contain the appropriate functional group(s) necessary to form an indole ring substituted by  $R_1$ ,  $R_2$  and  $R_3$  as defined in claim 1 or groups convertible thereto.

36. A pharmaceutical composition which comprises a compound according to claim 1 and a pharmaceutically acceptable carrier.

10

37. A compound according to claim 1 for use as a therapeutic substance.

38. A compound according to claim 1 for use in the treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse and/or schizophrenia.


39. A method of treatment or prophylaxis of anxiety, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse and/or schizophrenia in mammals including humans, which comprises administering to the sufferer a therapeutically effective amount of a compound according to claim 1.

40. The use of a compound according to claim 1 in the manufacture of a medicament for the treatment or prophylaxis of anxiety, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse and/or schizophrenia.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 91/01553

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (if several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C07D401/12; A61K31/44		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.Cl. 5	C07D	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>10</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
A	JOURNAL OF MEDICINAL CHEMISTRY. vol. 29, no. 11, November 1986, WASHINGTON US pages 2415 - 2418; P. FLUDZINSKI: '2,3-dialkyl(dimethylamino)indoles: Interaction with 5HT <sub>1</sub> , 5HT <sub>2</sub> and rat stomach fundal serotonin receptors.' cited in the application see the whole document  ---	1, 36, 37
<p><sup>10</sup> Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
05 DECEMBER 1991	16. 12. 91	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	B. S. de Jong 	

GB 9101553  
SA 51152

**This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on**  
**The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 05/12/91**

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
/*- -		FR-A- 1515687 GB-A- 115410 NL-C- 133871 NL-A- 660376	
<hr/>			